

Abstract

Analysis and Usage Perspective of Solid Digestate [†]

Eglė Didžiulytė * and Rasa Šlinkšienė 

Department of Physical and Inorganic Chemistry, Kaunas University of Technology, LT-50254 Kaunas, Lithuania; rasa.slinksiene@ktu.lt

* Correspondence: egle.didziulyte@ktu.lt

[†] Presented at the International Conference EcoBalt 2023 “Chemicals & Environment”, Tallinn, Estonia, 9–11 October 2023.

Keywords: digestate; organic farming; organic carbon

Extremely intensive and chemicalized agriculture severely damages the naturally formed eco-system [1]. This issue encourages concern about soil conservation and improvement of its quality, human health, and environmental protection; therefore, it is necessary to look for alternative methods of agriculture. One of these methods is organic farming, where mineral fertilizers are not used and nutrients needed for plants are provided using biological substances of organic origin. On the other hand, the accumulation of such biodegradable organic waste and its disposal in landfills causes various environmental, economic, and social problems. One such waste is digestate [2]. Most of the digestate (about 120 million tons) is agricultural digestate (a mixture of manure and plants). The rest is obtained by mechanical biological treatment from municipal solid waste, separated biological waste, sewage sludge, or the agriculture and food industries. In various EU countries, renewable energy policies and subsidies for the production of electricity, gas, and heat from biomass have improved economic conditions for the anaerobic digestion of biological waste or food waste, but in this process, digestate is also produced. However, it was also noted that a lot of detailed scientific research should be conducted concerning the risk factors, environmental effects, fertilizer treatment methods, and enrichment during long-term use [3–5]. Because the digestate is rich in organic matter, during this study, when three different digestates were used, the organic carbon content was first determined. The tests were carried out on dried digestates, which were of two types: whole and chopped. It was found that the content of organic carbon is higher in whole samples and ranges from 39.0 to 50.0%, and ranges from 34.8 to 45.2% in chopped. In order to obtain the maximum concentration of organic carbon from the used digestates, extracts were produced on the basis of water, potassium alkali, and different production conditions. The samples were kept for 3, 6, and 9 days at room temperature and for 3, 6, and 9 h at different temperatures, mixed with a mechanical stirrer, and in an ultrasonic bath. Also, the samples were heated at temperatures of 50 °C, 70 °C, and 90 °C. A higher concentration of organic carbon was found in potassium-based extracts, and the higher temperature had a positive effect on the concentration of organic carbon in the samples.



Citation: Didžiulytė, E.; Šlinkšienė, R. Analysis and Usage Perspective of Solid Digestate. *Proceedings* **2023**, *92*, 34. <https://doi.org/10.3390/proceedings2023092034>

Academic Editors: Monika Mortimer, Anne Kahru, Ivo Leito, Riin Rebane and Villem Aruoja

Published: 22 November 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Author Contributions: Conceptualization, R.Š.; methodology, E.D. and R.Š.; software, E.D.; resources, E.D.; data curation, R.Š. and E.D.; writing—original draft preparation, E.D.; writing—review and editing, R.Š. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data sharing not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Parikh, S.J.; James, B.R. Soil: The Foundation of Agriculture. *Nat. Educ. Knowl.* **2012**, *3*, 2.
2. Paull, J. Attending the First Organic Agriculture Course: Rudolf Steiner's Agriculture Course at Koberwitz, 1924. *Eur. J. Soc. Sci.* **2011**, *21*, 64–70.
3. Kovačić, Đ.; Lončarić, Z.; Jović, J.; Samac, D.; Popović, B.; Tišma, M. Digestate Management and Processing Practices: A Review. *Appl. Sci.* **2022**, *12*, 9216. [[CrossRef](#)]
4. Cheffins, N.; Stainton, D. *Final Project Report for OMK006-06*; WRAP: Banbury, UK, 2015.
5. Baştalık, B.; Koçar, G. A review of the biogas digestate in agricultural framework. *J. Mater. Cycles Waste Manag.* **2020**, *22*, 1318–1327. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.