

Exploring CO₂ storage potential in Lithuanian deep saline aquifers using digital rock volumes: a machine learning guided approach

Shruti Malik¹, Pijus Makauskas², Ravi Sharma³, Mayur Pal⁴

^{1,2,4}Kaunas University of Technology, Department of Mathematical Modelling, Kaunas, Lithuania

³Department of Earth Sciences, Indian Institute of Technology Roorkee, India

⁴Corresponding author

E-mail: ¹shruti.malik@ktu.lt, ²pijus.makauskas@ktu.lt, ³ravi.sharma@es.iitr.ac.in, ⁴mayur.pal@ktu.lt

Received 30 November 2023; accepted 28 December 2023; published online 31 December 2023

DOI <https://doi.org/10.21595/accus.2023.23906>



Copyright © 2023 Shruti Malik, et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract. The increasing significance of carbon capture, utilization and storage (CCUS) as a climate mitigation strategy has underscored the importance of accurately evaluating subsurface reservoirs for CO₂ sequestration. In this context, digital rock volumes, obtained through advanced imaging techniques such as micro-Xray computed tomography (MXCT), offer intricate insights into the porous and permeable structures of geological formations. This study presents a comprehensive methodology for assessing CO₂ storage viability within Lithuanian deep saline aquifers, namely Syderiai and Vaskai, by utilizing petrophysical properties estimated from digital rock volumes of samples from analogous formations. It also demonstrates the potential of integrating advanced imaging techniques, machine learning, and numerical modeling for accurate assessment and effective management of subsurface CO₂ storage.

Keywords: carbon capture, utilization and storage (CCUS), saline aquifers, storage potential, digital rock volumes, machine learning, lattice Boltzmann method, numerical modeling.

1. Introduction

In recent years, anthropogenic activities have led to a global increase in greenhouse gas emissions. To mitigate this, Carbon Capture, Utilization, and Storage (CCUS) has emerged as a potential solution [1]. In Lithuania's Baltic Basin, research is still in its early stages regarding the long-term fate of geological CO₂ storage [2, 3]. This study focuses on the deep saline aquifers, Syderiai and Vaskai of the Baltic Basin (shown in Fig. 1) and aims to demonstrate the effective application of machine learning in extracting optimized estimates of storage and flow potential using non-destructive digital rock volumes (DRV).




Fig. 1. Location of Syderiai and Vaskai regions

Machine learning algorithms offer an accurate and fast alternative to time-consuming conventional Digital rock physics method for determining optimized estimates of petrophysical properties [4-6]. The storage of captured CO₂ can be done in underground geological formations such as depleted oil and gas reservoirs, deep saline aquifers, or coal seams. Amongst these, deep saline aquifers are considered as the most prospective site due to their large storage capacity and widespread geographic distribution, making it easier to find storage locations closer to the sources of CO₂ emissions [3].

2. Methodology

In this study, digital volumes of rocks were obtained from formations analogous to Lithuanian reservoirs using Micro Xray Computed Tomography (MXCT) scanning technique. The Machine learning (ML) algorithm was then employed to estimate the porosity, and Lattice Boltzmann Method (LBM) simulations were conducted to estimate permeability values of 3D DRVs [7]. Sub-volumes were extracted from segmented volumes to investigate fluid flow behavior and determine the representative element volumes (REV) (as shown in Fig. 2). These sub-volumes can also be used to analyze the geo-chemical aspect which includes the impact of fluids on porosity and its distribution.




Fig. 2. Illustration of the sub-volume extraction for REV estimation

3. Results

The estimated petrophysical properties were compared with those measured in laboratory. It was observed that the estimated values closely matched the laboratory measurements. The error percentages for porosity were in the range of ‘1 %-8 %’, while for permeability (Table 1) they ranged from ‘20 %-55 %’, as shown in Table 1.

Table 1. Permeability estimation on sub-volumes of samples from analogous formation using LBM

Sample	Average permeability of sub-volumes (mD)	Laboratory (mD)	Error (%)
S1	330	275	20
S2	96	62	55
S3	396	327	21




Fig. 3. Mechanistic model showing mid-case permeability distribution for Vaskai aquifer [8]

Further, the numerical modeling of CO₂ injection into saline aquifers was performed, to estimate the storage capacity using tNavigator software [8, 9]. Three test cases were defined, wherein the reservoir properties extracted from published literature served as the mid-case values. A low case was defined, in which parameters were decreased by 10 % from the mid-case values, and a high case where the parameters were increased by 30 % from the mid-case values [8]. The mechanistic model showing the mid-case permeability distribution for Vaskai aquifer is shown in Fig. 3.

4. Conclusions

The current study aims to explore the CO₂ storage potential of deep saline aquifers in Lithuania, specifically the Syderiai and Vaskai formations, using numerical modeling of CO₂ injection into these formations. Measurements on samples from formations analogous to those in Lithuania were used to establish a baseline understanding, which will aid in the analysis of Lithuanian reservoir samples.

Further, this work will be extended to include samples from actual Lithuanian reservoirs and to study the geochemical reactions and geo-mechanical behaviour of the rocks. Such studies shall further facilitate identification of reservoir(s) wherein sequestration potential can be reliably explored.

Acknowledgements

The authors would like to acknowledge the Lithuanian Research Council Funding for postdoctoral research fund proposal registration No. P-PD-22-022-PATIKSLINTA, the support from UAB Minijos Nafta for sharing data for reservoir modeling and simulation, and Rock Flow Dynamics for sharing Educational License of T-navigator for simulation performed in this study.

Data availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

Author contributions

Shruti Malik: investigation, methodology, software, writing – original draft preparation. Pijus Makauskas: methodology, software, writing – review and editing. Ravi Sharma: methodology, supervision, writing – review and editing. Mayur Pal: conceptualization, investigation, methodology, supervision, writing – review and editing.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- [1] A. Shogenova, S. Šliaupa, K. Shogenov, R. Vaher, and R. Šliaupienė, “Geological Storage of CO₂ – Prospects in the Baltic States,” in *69th EAGE Conference and Exhibition incorporating SPE EUROPEC 2007*, Jan. 2007, <https://doi.org/10.3997/2214-4609.201401887>
- [2] R. Šliaupienė and S. Šliaupa, “Prospects for CO₂ geological storage in deep saline aquifers of Lithuania and adjacent territories,” *Geologija*, Vol. 53, No. 3(75), pp. 121–133, 2011.
- [3] S. Šliaupa and R. Šliaupienė, “Prospects of Geological Storage of CO₂ in Lithuania,” in *Baltic Carbon Forum*, Oct. 2021.

- [4] N. Alqahtani, F. Alzubaidi, R. T. Armstrong, P. Swietojanski, and P. Mostaghimi, “Machine learning for predicting properties of porous media from 2d X-ray images,” *Journal of Petroleum Science and Engineering*, Vol. 184, p. 106514, Jan. 2020, <https://doi.org/10.1016/j.petrol.2019.106514>
- [5] D. Tang and K. Spikes, “Segmentation of shale SEM images using machine learning,” in *SEG Technical Program Expanded Abstracts 2017*, Aug. 2017, <https://doi.org/10.1190/segam2017-17738502.1>
- [6] Y. D. Wang, M. J. Blunt, R. T. Armstrong, and P. Mostaghimi, “Deep learning in pore scale imaging and modeling,” *Earth-Science Reviews*, Vol. 215, p. 103555, Apr. 2021, <https://doi.org/10.1016/j.earscirev.2021.103555>
- [7] S. Malik, P. Makauskas, V. Karaliute, R. Sharma, and M. Pal, “Assessing Long-term fate of geological CO₂ storage in Lithuania: A machine learning approach for pore-scale processes and reservoir characterization,” in *12th Trondheim Conference on CO₂ Capture, Transport and Storage*, 2023.
- [8] M. Pal, S. Malik, V. Karaliūtė, P. Makauskas, and R. Sharma, “assessing the feasibility of carbon capture and storage potential in Lithuanian geological formations: a simulation-based assessment,” in *84th EAGE Annual Conference and Exhibition*, Vol. 2023, No. 1, pp. 1–5, Jan. 2023, <https://doi.org/10.3997/2214-4609.202310502>
- [9] “T-Navigator, Reservoir Simulation Software Version 21.3, Rock flow Dynamics,” 2022.



Dr. **Shruti Malik** earned her Ph.D. degree from Indian Institute of Technology, Roorkee, India. She is working as a post-doctoral researcher in the Department of Mathematical Modelling, Kaunas University of Technology (KTU), Lithuania, supported by a research grant from Research Council of Lithuania. Her work is focused on blending the digital rock physics with AI & ML techniques to assess the impact of CO₂ & H₂ storage on the sub-surface reservoirs.



Pijus Makauskas is a Ph.D. researcher at the department of Mathematical Modelling in Faculty of Mathematics and Natural sciences at Kaunas University of Technology. His area of research includes subsurface modelling, Machine learning and AI, Geothermal, CO₂ and Hydrogen storage, numerical modelling of subsurface flows and reservoir simulation.



Dr. **Ravi Sharma** received a master’s degree in applied Geophysics from the University of Roorkee in 1999. He received his MS and Ph.D. in Petroleum Engineering from Colorado School of Mines, USA, in 2015. He has extensive work experience in various roles with the hydrocarbon energy industry. His research interest includes experimental and modelling methods in rock physics and petrophysics for storage, flow, and the associated elastic and geo-mechanical property determination, integrated reservoir (convention and unconventional) characterization, ML & AI applications in geosciences and petroleum engineering. Dr. Sharma is an active member of AGU, IEEE, SEG, SPE, AAPG, SPG, and SPWLA, an Associate Editor of the Journal of Applied Geophysics and Geohorizons, and a guest editor of Frontiers in Earth science.



Dr. **Mayur Pal** received a Ph.D. degree in computational engineering from Swansea University, Wales, UK in 2007. After finishing his Ph.D. Mayur Pal worked at Shell International Exploration and Production B.V. Research Centre in Rijswijk followed by Maersk Oil Research and Technology Centre. He also held positions as head of enhanced oil recovery team and head of asset in North Oil Company, Qatar. He is currently Prof. at Department of Mathematical Modelling at KTU, Kaunas, Lithuania. His research interest includes, subsurface flows, multiscale modelling, discrete fracture network modelling, enhanced oil recovery, CCUS, data science and machine learning applications to solve problems in engineering and sciences.